

Clinical features of anaphylaxis according to age in a single university hospital in Korea

Lira Yoon,^{1,2} Bo Ryung Kim,^{1,2} Jae Young Lee,³ Kwanghoon Kim,^{1,2} Young Mi Kim,^{1,2} Seong Heon Kim,^{4,5} Hye-young Kim^{1,2}

Abstract

Background: Prevalence of anaphylaxis is increasing steadily, but there is scant information about anaphylaxis.

Objective: To identify the prevalence of anaphylaxis and evaluate the cause and features of anaphylaxis according to age.

Method: The relevance of gender, age, history of allergic diseases, causes and clinical symptoms in the treatment and prognosis of anaphylaxis were retrospectively evaluated in patients diagnosed with anaphylaxis in Pusan National University Hospital from January 2009 to June 2014.

Results: The 161 patients included 30 children and 131 adults. The prevalence of anaphylaxis in both children and adults increased during the study period. The most common triggers were food in children, especially in those <5 years of age, and drugs in adults. All of the children with episodes of anaphylaxis presented with cutaneous symptoms and 90% had respiratory symptoms. For adults, cardiovascular symptoms were more common.

Conclusions: The prevalence of anaphylaxis increased in both children and adults, with a higher rate in children. The causes and symptoms of anaphylaxis may differ depending on age. Clinicians should understand the characteristics of anaphylaxis by age.

Keywords: Anaphylaxis, age, epidemiology, etiology, children, infant

From:

¹ Department of Pediatrics, Pusan National University School of Medicine

² Medical Research Institute of Pusan National University Hospital, Busan, Korea

³ Department of nursing science, Kyungsoo University, Busan, Korea

⁴ Department of Pediatrics, Pusan National University Children's Hospital, Yangsan, Korea

⁵ Research Institute for Convergence of Biomedical Science and Technology, Pusan National University Yangsan Hospital

Corresponding author:

1. Hye-young Kim

Department of Pediatrics, Pusan National University Hospital, Pusan National University

College of Medicine, 179 Guduck-ro, Seo-gu, Busan, Korea

E-mail: pearlhy@naver.com

2. Seong Heon Kim

Department of Pediatrics Pusan National University Children's

Hospital, Pusan National University College of Medicine, Kumoh-ro 20, Mulgum-up, Yangsan, Gyeongnam, Korea

E-mail: pedksh@gmail.com

Introduction

Anaphylaxis is a systemic allergic reaction with a rapidly progressive onset that can be fatal. It requires immediate diagnosis and treatment.¹⁻³ The incidence and severity of allergic diseases is steadily increasing in Korea, causing an intense awareness and an interest in anaphylaxis.⁴⁻⁷ It is important to understand the epidemiology of anaphylaxis in order to manage patients properly and prevent recurrence. However, the range of prevalence and triggering factors of anaphylaxis varies worldwide.⁷⁻¹⁰ The epidemiology of anaphylaxis in Asia is unclear.

The incidence of anaphylaxis in Korea, as reported by Lim¹¹

by analyzing the medical claim data of the Health Insurance Review and Assessment Service (HIRA) from 2001 to 2007 under the disease code 'anaphylaxis', was 0.7-1.0 cases per 100,000 persons under 18 years of age. Lee et al.⁴ analyzed patients under 18 years of age who visited one of nine general hospital emergency rooms between June 2008 and March 2009 (10 months), and reported that 7.23 cases out of 10,000 pediatric patients were diagnosed with anaphylaxis. Yang et al. reported that among 978,146 patients, 138 (0.014%) patients had anaphylaxis in a single center.¹² However, there is no comprehensive research on the incidence and prevalence of anaphylaxis in Korea.

As anaphylaxis shows a variety of clinical symptoms and can be life threatening, many guidelines have been proposed for a rapid diagnosis and effective treatment; however, the diagnostic rate and level of epinephrine prescription or self-administration is still reported to be low.¹³ In particular, children have difficulties expressing their symptoms. The younger the patient is, the harder it is to diagnose and treat them.³ Also, there have been many studies on the characteristics of anaphylaxis in adults according to age, so efforts are needed to comprehend the characteristics of anaphylaxis by age. We analyzed the incidence of patients diagnosed with anaphylaxis in our hospital during 5 years and aimed to determine the cause and clinical characteristics of anaphylaxis according to age.

Methods

A retrospective chart review was performed in anaphylaxis patients at Pusan National University Hospital, Busan, Korea from January 2009 to June 2014. Our university hospital is a reference center for a population of 6 million inhabitants in Busan and Gyeongnam. Information on sex, age, history of allergic diseases, cause of anaphylaxis, clinical symptoms, treatment, and prognosis were collected. Subjects under 19 years of age were defined as children, and those above 20 years of age were defined as adults. All patients were further classified into six age groups: under 5 years, 6-11 years, 12-19 years, 20-39 years, 40-59 years, and more than 60 years. Patients were first selected using the disease codes for anaphylaxis, anaphylaxis shock, allergy and anaphylaxis, hypersensitivity, food hypersensitivity, and drug hypersensitivity on the basis of the International Statistical Classification of Diseases, 10th Revision (ICD-10), and their medical records were analyzed so that only patients full filling the diagnostic criteria for anaphylaxis were included. Anaphylaxis was diagnosed if one of the following three criteria were met: the occurrence of respiratory or cardiovascular symptoms accompanied by an acute cutaneous reaction within a few minutes to a few hours; the occurrence of two or more systemic responses out of the

cutaneous, respiratory, cardiovascular, or gastrointestinal system, after exposure to an allergic antigen; and the occurrence of hypotension after exposure to a known allergic antigen.¹ The study was approved by the Pusan National University Hospital Institutional Review Board.

Statistical analyses were performed using IBM SPSS ver. 18.0 (IBM Co., Armonk, NY, USA), and the Chi-square test and Fisher's exact test were used to test the statistical significance in children and adults. Data were considered to be statistically significant if the p-value was < 0.05.

Results

There were 161 subjects included in the study, with 30 children and 131 adults. Sixteen (53.3%) were male children and 65 (49.6%) were male adults. The average age at diagnosis was 9.5 ± 5.7 years for children and 50.2 ± 16.0 years for adults (Table 1). There were 22 children (73.3%) and 69 adults (52.7%)

Table 1. Demographic characteristics of patients with anaphylaxis

	Children (n=30)	Adult (n=131)	P-value
Sex			
Male	16(53.3)	65(49.6)	
Female	14(46.7)	66(50.4)	
Mean Age (yr)	9.5±5.7	50.2±16.0	
History of allergic diseases			
Atopic dermatitis	13(43.3)	2(1.5)	0.000*
Food allergy	13(43.3)	28(21.4)	0.019*
Allergy rhinitis	10(33.3)	9(6.9)	0.000*
Asthma	8(26.7)	12(9.2)	0.014*
Chronic urticaria	2(6.7)	12(9.2)	1.000
Drug allergy	1(3.3)	24(18.3)	0.049*

Values are presented as mean±standard deviation (range), number (%).
*P<.05 was considered statistically significant.




Figure 1. Incidence of anaphylaxis in Pusan National University Hospital from 1.7 per 100,000 patients in 2009 to 60.2 per 100,000 patients in 2014 in children and 0.9 per 100,000 patients in 2009 to 14.7 per 100,000 patients in 2014 in adult group.

with a history of allergic disease (Table 1). Histories of atopic dermatitis, food allergy, allergic rhinitis, and asthma were higher in children, while drug allergy was higher in adults.

The incidence of anaphylaxis increased during the study period in both children and adults (Figure 1). In children, the number of patients who visited the hospital increased steadily from 1.7 per 100,000 in 2009, 3.6 in 2010, 11.9 in 2011, 13.3 in 2012, and 30.9 in 2013, to 60.2 in 2014. For adults, the number of patients who visited the hospital also increased from 0.9 per 100,000 in 2009 to 14.7 in 2014.

Causes of anaphylaxis in children included food (50%), drugs (26.7%), food-dependent exercise-induced anaphylaxis (6.7%), blood transfusion (3.3%), insect bites (3.3%), and idiopathic causes (10%). Causes in adults included drugs (55.7%), food (22.1%), insect bites (6.9%), exercise (4.6%), food-dependent exercise-induced anaphylaxis (3.8%), and idiopathic causes (6.9%) (Table 2). Among the food allergens, milk and seafood were the most common causes in children and adults, respectively, and cephalosporin and non-steroidal anti-inflammatory drugs (NSAIDs) were the most common causative drugs for anaphylaxis (Table 2). Anaphylaxis due to food was the most common in children under 5 years old, at 88.9% (Table 4).

Cutaneous symptoms occurred in 30 children (100%) and 98 adults (74.8%), respiratory symptoms occurred in 27

Table 2. Underlying causes of anaphylaxis

Underlying cause	Children (n=30)	Adult (n=131)	P-value
Food	15(50.0)	29(22.1)	0.003*
Cow's milk	5(16.7)	0	
Egg white	2(6.7)	0	
Sea foods	1(3.3)	9(6.9)	
Fruits/vegetables	1(3.3)	5(3.8)	
Wheat	1(3.3)	7(5.3)	
Nuts	0	3(2.3)	
Soy	1(3.3)	2(1.5)	
Buckwheat	1(3.3)	1(0.8)	
Chicken	1(3.3)	0	
Beef	1(3.3)	1(0.8)	
Pork	1(3.3)	1(0.8)	
Drug	8(26.7)	73(55.7)	0.005*
IgE mediated			
Ampicillin	0	3(2.3)	
Cephalosporin	3(10.0)	22(16.8)	
Muscle relaxant	0	3(2.3)	
Methylprednisolone	1(3.3)	1(0.8)	
Non-IgE mediated			
Aspirin or NSAID	4(13.3)	23(17.6)	
Radio contrast	0	8(6.1)	
Acetaminophen	0	3(2.3)	
Other	0	10(7.6)	
Exercise(food dependent)	2(6.7)	5(3.8)	0.615
Exercise(food independent)	0	6(4.6)	0.594
Insect sting	1(3.3)	9(6.9)	0.690
Blood transfusion	1(3.3)	0	0.186
Idiopathic	3(10.0)	9(6.9)	0.698

Values are presented as number (%).

*P<.05 was considered statistically significant.

children (90%) and 82 adults (62.6%), gastrointestinal symptoms occurred in 14 children (46.7%) and 36 adults (27.5%), and cardiovascular symptoms occurred in 8 children (26.7%) and 81 adults (61.8%). Cutaneous, respiratory, and gastrointestinal symptoms were significantly more common in children compared to adults (Table 3). Cardiovascular symptoms were more common in adults compared to children and were not seen in children under 5 years of age and increased in proportion to age (Table 4). A biphasic response was seen in 7 children (23.3%) and 3 adults (2.3%), which was significantly more common in children (Table 3). The time interval from exposure to a causal agent to the appearance of symptoms was within 10 minutes in 10 children (33.3%), under 30 minutes in 8 children (26.7%), within 10 minutes in 32 adults (24.4%), and under 30 minutes in 37 adults (28.2%).

Twenty-two children (73.3%) and 27 adults (20.6%) received epinephrine treatment, and 12 children (40.0%) and 23 adults (17.6%) were prescribed a prophylactic self-injectable epinephrine (Table 5). In children, the younger they were, the less likely they received epinephrine and prophylactic self-injectable epinephrine prescription; no difference was seen in adult subjects (Table 4). Seven adults (6.8%) received

Table 3. Signs and symptoms of anaphylaxis

	Children (n=30)	Adult (n=131)	P-value
Cutaneous	30(100.0)	98(74.8)	0.001*
Urticaria	25(83.3)	58(44.3)	
Itch-flush	22(73.3)	42(32.1)	
Pruritus and swelling of lips-tongue-uvula	7(23.3)	6(4.6)	
Erythema	7(23.3)	18(13.7)	
Angioedema	6(20.0)	21(16.0)	
Respiratory	27(90.0)	82(62.6)	0.004*
Dyspnea	23(76.7)	76(58.0)	
Cough	14(46.7)	5(3.8)	
Wheeze	10(33.3)	7(5.3)	
Stridor	9(30.0)	7(5.3)	
Coryza	5(16.7)	0	
Cyanosis	5(16.7)	13(9.9)	
Cardiovascular	8(26.7)	81(61.8)	0.000*
Hypotension	7(23.3)	70(53.4)	
Pallor	5(16.7)	0	
Chest pain	2(6.7)	8(6.1)	
Collapse	1(3.3)	1(0.8)	
Diaphoresis	0	8(6.1)	
Cardiac arrest	0	7(5.3)	
Gastrointestinal	14(46.7)	36(27.5)	0.049*
Nausea	11(36.7)	14(10.7)	
Vomiting	10(33.3)	17(13.0)	
Abdominal pain	4(13.3)	17(13.0)	
Neurologic	14(46.7)	73(55.7)	0.420
Dizziness	8(26.7)	32(24.4)	
Anxiety	5(16.7)	1(0.8)	
Confusion	1(3.3)	3(2.3)	
Loss of consciousness	1(3.3)	44(33.6)	
Death	0	1(0.8)	
Biphasic response	7(23.3)	3(2.3)	0.000*

Values are presented as number (%).

*P<.05 was considered statistically significant.

Table 4. Underlying cause, signs, symptoms and treatment according to age group

	Age groups					
	≤5yr	6-11yr	12-19yr	20-39yr	40-59yr	≥ 60yr
Number	9(5.6)	9(5.6)	12(7.5)	37(23.0)	56(34.8)	38(23.6)
Cause						
Food	8(88.9)	3(33.3)	4(33.3)	9(24.3)	10(17.9)	10(26.3)
Drug	1(11.1)	3(33.3)	5(41.7)	19(51.4)	32(57.1)	22(57.9)
Exercise(food)	0	0	2(16.7)	3(8.1)	2(3.6)	0
Exercise	0	0	0	2(5.4)	4(7.1)	0
Insect sting	0	1(11.1)	0	1(2.7)	6(10.7)	2(5.3)
Idiopathic	0	2(22.2)	1(8.3)	3(8.1)	2(3.6)	4(10.5)
Signs and symptoms						
Cutaneous	9(100.0)	9(100.0)	12(100.0)	32(86.5)	40(71.4)	26(68.4)
Respiratory	9(100.0)	9(100.0)	9(75.0)	28(75.7)	34(60.7)	20(52.6)
Cardiovascular	0	3(33.3)	5(41.6)	18(48.6)	34(60.7)	29(76.3)
Gastrointestinal	4(44.4)	4(44.4)	4(33.3)	11(29.7)	19(33.9)	6(15.8)
Neurological	4(44.4)	4(44.4)	6(50.0)	10(27.0)	36(64.3)	27(71.1)
Treatment						
Epinephrine	4(44.4)	9(100.0)	9(75.0)	4(10.8)	15(26.8)	8(21.1)
Prophylactic self-injectable epinephrine	3(33.3)	4(44.4)	5(41.7)	6(16.2)	10(17.9)	7(18.4)

Values are presented as number (%).

Table 5. Treatment of anaphylaxis

	Children (n=30)	Adult (n=131)
Emergency treatment		
Epinephrine	22(73.3)	27(20.6)
Antihistamine	20(66.7)	63(48.1)
Systemic steroid	18(60.0)	62(47.3)
IV hydration	18(60.0)	44(33.6)
Bronchodilator	12(40.0)	10(7.6)
O2 inhalation	13(43.3)	43(32.8)
Prophylactic self-injectable epinephrine	12(40.0)	23(17.6)
Hospitalization	17(56.7)	16(12.2)

Values are presented as number (%).

treatment in the intensive care unit, 7 (6.8%) were hospitalized, and there was one death from a bee sting-induced anaphylaxis. One child (3.3%) received treatment in the intensive care unit and 16 (53.3%) were hospitalized, with no cases of death reported.

Discussion

The lifetime prevalence of anaphylaxis is about 0.05-2%,^{2,14,15} and clinical studies show that the incidence is steadily increasing.^{3,12,16-21} In this study, the incidence of anaphylaxis for last 5 years also increased, especially at a higher rate in children. Recent nationwide data in Korean adults also showed that the incidence of anaphylaxis significantly increased from 7.74 cases per 100,000 in 2007 to 13.32 in 2011.⁷ This increase in incidence is thought to be due to an increased diagnostic rate following the setup of guidelines, and also due to the increased prevalence of allergic diseases in general.^{2,20,22} In our study, the reason for the increase in pediatric anaphylaxis incidence being even more prominent may be because of the active education and

promotion of anaphylaxis following the initiation of pediatric allergist specialist clinics in 2009. However, more focus and continuous research is needed on the rapid increase of anaphylaxis incidence in children in this study.

In this study, the most common cause of anaphylaxis in children was food, especially in children under 5 years old. Although food is known a common cause of pediatric anaphylaxis, the type of food has differed between studies.^{3,4,12,16,19,23-25} In Western studies including the United States, peanuts and other nuts are the most common causes.^{3,16,20,23,25-27} In Korea, the most commonly reported causes have been buckwheat,¹¹ fish,⁴ and eggs.²⁸ In this study, milk was found to be the most common cause. Such differences are generally thought to be due to the inclusion criteria of the subjects or to differences in the age or region of the subject.^{18,20,29}

In the present study, as in other similar studies,^{12,30-33} drugs, food, and insect stings were the most common causes of anaphylaxis in adults, with no difference according to age. Among the drugs, NSAIDs, and cephalosporin were most common. Seafood was the most common food to cause anaphylaxis in adults, in agreement with the results of a study on food anaphylaxis in adults living in the Southeastern seaside,¹⁷ which can be presumed to be a characteristic of Korean adults living close to the shoreline.^{30,34} In this study, the incidence of anaphylaxis due to insect stings is lower than in other studies.^{7,12} We suggest the reason for this being that only some of the patients referred to our university hospital resided in an urban area, because insect stings often occur in rural settings and cause mild anaphylaxis symptoms.⁷

Presently, 73.3% of children and 52.7% of adults had a history of allergic disease. The history of atopic dermatitis, food allergy, allergic rhinitis, and asthma was higher in children, while drug allergy was higher in adults. In other studies, allergy history was reported to be higher in children than adults, with

prevalence rates of 64 to 82% in children^{22,31,33} and 19 to 50% in adults.^{7,10,12} These differences may reflect the increasing prevalence of allergy disease. As the hospitalization rate of children was higher than adults in our study, it might be possible to obtain a more detailed medical history.

The most common clinical manifestation of anaphylaxis has been reported to be skin reactions,^{4,12,20,22,24,25,28-35} which is also the case in this study for both children and adults. Cutaneous, respiratory, and gastrointestinal symptoms were more commonly observed in children compared to adults, and cardiovascular symptoms were more common in adults. In particular, cardiovascular symptoms were not seen in children under 5 years of age and increased with age. It is hard to measure blood pressure in children and tachycardia is a more common accompanying symptom rather than hypotension, so caution is needed for diagnosis.^{3,24,25} Also, the reason why cardiovascular symptoms are more common in adults may be because drugs are more frequently involved as a cause, and there are more patients with a history of cardiovascular disease as their age increases.^{24,35} In contrast to previous reports that biphasic responses are uncommon in children,^{20,31-33,36} they were more common in children in this study. These results may be influenced by the fact that there were more cases of hospitalization in children, so they were observed for a longer period of time.

In this study, the prescription rate of epinephrine and portable epinephrine was higher in children compared to adults. However, the prescription rate under 5 years of age was lower compared to the other pediatric age groups. This is because subjective symptoms are ambiguous and blood pressure measurement is difficult in this age group, meaning that diagnosis is not easy, and also because prophylactic self-injectable epinephrine is hard to use in children who weigh less than 10kg.

This study is limited as the sample size was small and was a retrospective study in a single university hospital. However, the study is relevant as the increasing incidence of anaphylaxis was found and the characteristics of anaphylaxis were analyzed by age. Clinicians should understand the characteristics of anaphylaxis by age and try to diagnose actively and prescribe epinephrine. Also, the development of a prophylactic self-injectable epinephrine that can be used in children less than 10kg is necessary.

References

- Sampson HA, Munoz-Furlong A, Campbell RL, Adkinson NF, Jr., Bock SA, Branum A, et al. Second symposium on the definition and management of anaphylaxis: summary report--second National Institute of Allergy and Infectious Disease/Food Allergy and Anaphylaxis Network symposium. *Ann Emerg Med.* 2006;47:373-80.
- Jang GC, Chang Y-S, Choi SH, Song W-J, Lee S-Y, Park H-S, et al. Overview of anaphylaxis in Korea: diagnosis and management. *Allergy Asthma Respir Dis.* 2013;1:181-96.
- Simons FER, Sampson HA. Anaphylaxis: Unique aspects of clinical diagnosis and management in infants (birth to age 2 years). *J Allergy Clin Immunol.* 2015;135:1125-31.
- Lee SY, Kim KW, Lee HH, Lim DH, Chung HL, Kim SW, et al. Incidence and clinical characteristics of pediatric emergency department visits of children with severe food allergy. *Korean J Asthma Allergy Clin Immunol.* 2012;32:169-75.
- Gupta RS, Springston EE, Warriar MR, Smith B, Kumar R, Pongracic J, et al. The prevalence, severity, and distribution of childhood food allergy in the United States. *Pediatrics.* 2011;128:e9-e17.
- Amin AJ, Davis CM. Changes in prevalence and characteristics of IgE-mediated food allergies in children referred to a tertiary care center in 2003 and 2008. *Allergy and Asthma Proceedings: OceanSide Publications, Inc;* 2012. p. 95-101.
- Ye Y-M, Kim MK, Kang H-R, Kim T-B, Sohn S-W, Koh Y-I, et al. Predictors of the severity and serious outcomes of anaphylaxis in Korean adults: a multicenter retrospective case study. *Allergy Asthma Immunol Res.* 2015;7:22-9.
- Techapornroong M, Akrawinthawong K, Cheungpasitporn W, Ruxrungtham K. Anaphylaxis: a ten years inpatient retrospective study. *Asian Pac J Allergy Immunol.* 2010;28:262.
- Ben-Shoshan M, Clarke A. Anaphylaxis: past, present and future. *Allergy* 2011;66:1-14.
- Zubeldia J, Baeza M. Anaphylaxis in Adolescent/Adult Patients Treated in the Emergency Department: differences between initial impressions and the definitive diagnosis. *J Investig Allergol Clin Immunol.* 2015;25:288-94.
- Lim DH. Epidemiology of anaphylaxis in Korean children. *Korean J Pediatr.* 2008;51:351-4.
- Yang M-S, Lee S-H, Kim T-W, Kwon J-W, Lee S-M, Kim S-H, et al. Epidemiologic and clinical features of anaphylaxis in Korea. *Ann Allergy Asthma Immunol.* 2008;100:31-6.
- Simons FER, Arduzzo LR, Bilo MB, Dimov V, Ebisawa M, El-Gamal YM, et al. 2012 Update: World Allergy Organization Guidelines for the assessment and management of anaphylaxis. *Curr Opin Allergy Clin Immunol.* 2012;12:389-99.
- Frew A. What are the 'ideal' features of an adrenaline (epinephrine) auto-injector in the treatment of anaphylaxis? *Allergy.* 2011;66:15-24.
- Simons F, Arduzzo L, Bilò MB, El-Gamal YM, Ledford DK, Ring J, et al. World Allergy Organization guidelines for the assessment and management of anaphylaxis. *J Allergy Clin Immunol.* 2011;593:1-22.
- Altman AM, Camargo CA, Jr., Simons FE, Lieberman P, Sampson HA, Schwartz LB, et al. Anaphylaxis in America: A national physician survey. *J Allergy Clin Immunol.* 2015;135:830-3.
- Seo MH, Kim SH, Hong JS, Kim WY, Choe SW. Clinical features of food-induced anaphylaxis in the Southeastern coasted area of Korea. *Korean J Asthma Allergy Clin Immunol.* 2010;30:110-5.
- Wood RA, Camargo CA, Lieberman P, Sampson HA, Schwartz LB, Zitt M, et al. Anaphylaxis in America: the prevalence and characteristics of anaphylaxis in the United States. *J Allergy Clin Immunol.* 2014;133:461-7.
- Nocerino R, Leone L, Cosenza L, Canani RB. Increasing rate of hospitalizations for food-induced anaphylaxis in Italian children: An analysis of the Italian Ministry of Health database. *J Allergy Clin Immunol.* 2015;135:833.
- Chippis BE. Update in Pediatric Anaphylaxis A Systematic Review. *Clin Pediatr (Phila).* 2013;0009922812474683.
- Decker WW, Campbell RL, Manivannan V, Luke A, Sauver JLS, Weaver A, et al. The etiology and incidence of anaphylaxis in Rochester, Minnesota: a report from the Rochester Epidemiology Project. *J Allergy Clin Immunol.* 2008;122:1161-5.
- Russell S, Monroe K, Losek JD. Anaphylaxis management in the pediatric emergency department: opportunities for improvement. *Pediatr Emerg Care.* 2010;26:71-6.
- Muraro A, Roberts G, Clark A, Eigenmann PA, Halken S, Lack G, et al. The management of anaphylaxis in childhood: position paper of the European academy of allergology and clinical immunology. *Allergy.* 2007;62:857-71.
- Park HM, Noh JC, Park JH, Won YK, Hwang SH, Kim JY, et al. Clinical Features of Patients with Anaphylaxis at a Single Hospital. *Pediatr Allergy Respir Dis.* 2012;22:232.
- Rudders SA, Banerji A, Clark S, Camargo CA. Age-related differences in the clinical presentation of food-induced anaphylaxis. *J Pediatr.* 2011;158:326-8.
- Worm M, Eckermann O, Dölle S, Aberer W, Beyer K, Hawranek T, et al. Triggers and treatment of anaphylaxis: an analysis of 4000 cases from Germany, Austria and Switzerland. *Dtsch Arztebl Int.* 2014;111:367.
- Mehl A, Wahn U, Niggemann B. Anaphylactic reactions in children--a questionnaire-based survey in Germany. *Allergy.* 2005;60:1440-5.
- Roh EJ, Chung EH, Lee MH, Lee SJ, Youn YS, Lee JH, et al. Clinical features of anaphylaxis in the middle area of South Korea. *Pediatr Allergy Respir Dis.* 2008;18:61-9.
- Worm M, Moneret-Vautrin A, Scherer K, Lang R, Fernandez-Rivas M, Cardona V, et al. First European data from the network of severe allergic reactions (NORA). *Allergy.* 2014;69:1397-404.

30. Park HJ, Kim SH. Factors associated with shock in anaphylaxis. *Am J Emerg Med.* 2012;30:1674-8.
31. Orhan F, Canitez Y, Bakirtas A, Yilmaz O, Boz A, Can D, et al. Anaphylaxis in Turkish children: a multi-centre, retrospective, case study. *Clin Exp Allergy.* 2011;41:1767-76.
32. De Swert LF, Bullens D, Raes M, Dermaux AM. Anaphylaxis in referred pediatric patients: demographic and clinical features, triggers, and therapeutic approach. *Eur J Pediatr.* 2008;167:1251-61.
33. Hsin Y-C, Hsin Y-C, Huang J-L, Yeh K-W. Clinical features of adult and pediatric anaphylaxis in Taiwan. *Asian Pac J Allergy Immunol.* 2011;29:307-12.
34. Huang F, Chawla K, Jarvinen KM, Nowak-Wegrzyn A. Anaphylaxis in a New York City pediatric emergency department: triggers, treatments, and outcomes. *J Allergy Clin Immunol.* 2012;129:162-8 e1-3.
35. Braganza SC, Acworth JP, McKinnon DR, Peake JE, Brown AF. Paediatric emergency department anaphylaxis: different patterns from adults. *Arch Dis Child.* 2006;91:159-63.
36. Lee JM, Greenes DS. Biphasic anaphylactic reactions in pediatrics. *Pediatrics.* 2000;106:762-6.