

Erratum: Clinical features and predictive factors in neuropsychiatric lupus

Piyawan Chiewthanakul, Kittisak Sawanyawisuth, Chingching Foocharoen and Somsak Tiamkao

Dear Editor,

We are writing to you because we would like to do a minor change in an article entitled “Clinical features and predictive factors in neuropsychiatric lupus”, which has been published in Asian Pacific Journal of Allergy Immunology since March 2012 volume 30(1) page 55-60. Because there was a misunderstanding for the date of database collection, the duration of retrospective review should be 10 years starting from January 1996 to August 2006. We therefore would like to make an erratum for this article as below.

Abstract

From

Methods: Patients with SLE were retrospectively reviewed for 5 years, between January 2000 and August 2005. The prevalence, clinical features and predictive factors for NPL patients were studied. Neuropsychiatric (NP) syndromes were defined using the American College of Rheumatology (ACR) nomenclature and case definitions.

To

Methods: Patients with SLE were retrospectively reviewed for 10 years, between January 1996 and August 2005. The prevalence, clinical features and predictive factors for NPL patients were studied. Neuropsychiatric (NP) syndromes were defined using the American College of Rheumatology (ACR) nomenclature and case definitions.

Method

From

We enrolled 750 patients who were age over 15 years and met the diagnostic criteria of SLE5 in Srinagarind hospital, Khon Kaen, Thailand between January 2000 and August 2005.

To

We enrolled 750 patients who were age over 15 years and met the diagnostic criteria of SLE5 in Srinagarind hospital, Khon Kaen, Thailand between January 1996 and August 2005.

We apologize for making the errors. If you have any more questions, please let us know.

Kind regards,

Chingching Foocharoen

Department of Medicine, Faculty of Medicine,
Khon Kaen University, Khon Kaen, Thailand

