

Sputum Induction in Corticosteroid-Dependant Asthmatics: Risks and Airway Cellular Profile

L.C. Loh¹, V. Kanabar², M. D'Amato², N.C. Barnes³ and B.J. O'Connor²

SUMMARY Sputum induction with nebulized hypertonic saline is increasingly being used to evaluate airway inflammation. We investigated the procedure-associated risk in 16 asthmatics that were still symptomatic despite on high doses of regular corticosteroid (CS) therapy (7 on daily inhaled CS \geq 800 μ g budesonide or equivalent; 9 on additional daily oral CS) and their sputum cellular profile. For comparison, 12 mild stable asthmatics and 10 normal healthy subjects were included. All subjects inhaled 3%, 4% and 5% hypertonic saline sequentially via ultrasonic nebulizer as a means to induce sputum. Maximal percentage fall of Forced Expiratory Volume on One Second (FEV₁) during sputum induction was significantly greater in CS-dependent asthmatics (median % [IQR]: 16.0 [11.0-32.3]) than in mild asthmatics (5.3 [4.2-10.8], $p = 0.002$) and in normal subjects (4.6 [3.4-6.4]), $p = 0.0001$). The maximal percentage FEV₁ fall was inversely correlated with baseline FEV₁ ($R_s = -0.69$; $p < 0.0001$). Compared to mild asthmatics, induced sputum from CS-dependant asthmatics had proportionately fewer eosinophils (2.2 [0.8-7.0] versus 23.3% [10.7-46.3], $p = 0.003$) and greater neutrophils (64.2 [43.9-81.2] versus 28.7 [19.0-42.6], $p = 0.009$). Sputum neutrophils showed a significant inverse correlation to FEV₁ ($R_s = -0.51$, $p = 0.01$). We concluded that sputum induction using nebulized hypertonic saline should be performed with caution in CS-dependant asthmatics. The airway cellular profile observed suggests that the immunopathology underlying CS-dependant asthmatics may be different or a consequence of CS therapy.

The modern method of sputum induction is increasingly being used to evaluate airway inflammation in airway diseases.¹⁻³ With prior treatment with inhaled short-acting β_2 -agonist, nebulized hypertonic saline to induce sputum is generally shown to be safe and well tolerated in normal subjects and most asthmatic patients. Sputum induction has been attempted without serious hazards in adult asthmatic subjects recovering from severe exacerbations⁴ and in chronic stable asthmatics of greater severity.⁵⁻⁷ This relatively non-invasive approach to study air-

way inflammation is an invaluable alternative to bronchoscopy. This is particularly so in corticosteroid (CS)-dependant asthmatics whose easy susceptibility to bronchospasm precludes the use of bronchoscopy to investigate airway inflammation.

From the ¹Department of Medicine, International Medical University, Malaysia, ²Department of Respiratory Medicine and Allergy, Denmark Hill Campus, Guy's King's and St. Thomas's School of Medicine, London, ³Department of Respiratory Medicine, The London Chest Hospital, London, United Kingdom.
Correspondence: Li-Cher Loh
E-mail: loh@imu.edu.my

To date, the pathophysiology of airway inflammation in these asthmatics remains poorly understood. While eosinophils are widely regarded as the central effector cells in asthma pathogenesis,⁸⁻¹⁰ neutrophils had been shown to dominate in the airways of CS-dependent asthmatics^{7,11} compared to asthmatic patients of milder disease, suggesting that eosinophils may not be solely responsible in asthma pathogenesis.

In order to further explore the potential of sputum induction as a means to investigate airway inflammation, we studied the safety of sputum induction and the cellular profile of the induced sputum in symptomatic CS-dependant asthmatic patients, compared to those in mild stable asthmatic patients and in normal healthy subjects.

PATIENTS AND METHODS

Subjects

A group of asthmatic patients, being followed up in hospital chest clinics, was recruited according to eligibility as per study protocol. Normal healthy subjects were recruited by research advertisement. The diagnosis of asthma was defined according to international guidelines.¹² We defined 'CS-dependency' as dependency on inhaled budesonide ≥ 800 μg or equivalent daily with or without additional oral CS to provide best possible symptom control and 'symptomatic' as having ongoing asthmatic symptoms sufficient to interfere with daily living.¹³ These were individuals managed as \geq Step Four in the British Thoracic Society asthma guidelines¹⁴ with concomitant regular controller therapy such as theophylline, salmeterol or ipratropium bromide. Mild stable asthmatics were defined as having infrequent symptoms (i.e. \leq once weekly) that were easily relieved or prevented with rescue short-acting β_2 -agonist, and methacholine PC₂₀ of ≤ 8 mg/ml. Normal healthy subjects were defined as having no respiratory or other chronic medical conditions, non-smokers of at least 6 months duration and if previously smoked, had done so in less than 5 pack years. In addition, all normal subjects must demonstrate negative skin prick reactivity to cat, house dust mite or grass pollen, and methacholine PC₂₀ of ≥ 16 mg/ml. The study protocol was ap-

proved by local hospital ethics committees and informed written consent was obtained from all subjects.

Study design

All eligible subjects attended at least two visits: one screening and another where sputum induction was performed. During the screening visit, subjects' clinico-demographic details and lung functions were recorded. Normal subjects and mild asthmatics also had skin prick allergen test and methacholine inhalation challenge. Asthmatic patients were required to be clinically stable with unaltered regular asthma therapy for at least two weeks before the second visit. Where indicated, subjects were seen more than once in order to establish optimal treatment. Any subjects who had an exacerbation within three weeks would require that their second visit be postponed till a later date.

Sputum induction

We followed the method originally described by Pin and colleagues.¹⁵ Thirty minutes to one hour after treatment with salbutamol 200 μg via a meter dose inhaler (MDI) device (for mild asthmatics) or nebulized salbutamol 2.5 mg (for CS-dependant asthmatics), Forced Expiratory Volume in One Second (FEV₁) was recorded and this was taken as the baseline value for calculating the percentage FEV₁ change following hypertonic saline inhalation. Subjects then inhaled sequentially 3%, 4% and 5% hypertonic saline via an ultrasonic nebulizer, DeVilbiss Ultra-Neb 2000 (DeVilbiss Co, Heston, Middlesex, UK) in all subjects except oral CS-dependent asthmatics who were nebulized via Medix Sonic 2000 (Medix Ltd, Harlow, UK). This had occurred due to the availability of different ultrasonic nebulizers at the two different recruiting hospitals.

Each concentration of hypertonic saline was inhaled for 7 minutes, thereafter subjects rinsed their mouth and blew their nose (to minimize contamination from saliva or nose) before expectorating into a sterile container. FEV₁ was measured prior to each 7-minutes period and the procedure was terminated if FEV₁ fell below 20%. If the fall was between 10% and 20%, induction would be performed using the

same concentration of hypertonic saline as before (i.e. not to progress to higher concentration). A final FEV₁ was recorded and rescue salbutamol via nebulizer was given when clinically indicated.

Parameters for evaluating safety

The maximal percentage FEV₁ fall from baseline during the sputum induction, presence of hospital admission or prolonged observation more than one hour resulting from the sputum induction, were used as endpoints for evaluating safety.

Sputum processing and examination

The sputum was selected from saliva and processed within 2 hours according to a method described by Pizzichini and colleagues.¹ Briefly, sputum was homogenized by adding four volumes of freshly made 0.1% dithiothreitol (DTT) (Sputolysin, Calbiochem Ltd, Nottingham, UK) which was then added equal volume of Dulbecco phosphate-buffered saline (D-PBS) (Sigma, Poole, UK). The cell suspension was filtered through a 48 µm nylon gauze (BBSH Thompson, Scarborough, Ontario, Canada) and the filtrate centrifuged at 1,185 x g for 4 minutes at room temperature. The supernatant was aspirated and stored at -70°C for future assay while the cell pellet was resuspended with D-PBS, adjusted to 0.5 x 10⁶ cells/ml, to be placed into cups of Shandon III cytocentrifuge (Shandon, Inc. Pittsburgh, PA, USA) and cytospins made. After air dried, cytospins were stained with Wright's Giemsa for differential cell count on at least 400 non-squamous cells. These cytospins were counted by two independent observers. Cell counts were only accepted as valid if the differences of cell counts between the two observers were less than 5%. The average between the two counts was taken as the final count.

Data analysis

Descriptive statistics were used to summarize clinical characteristics of the study populations. Maximal percentage FEV₁ fall from sputum induction and the sputum cellular profile were expressed as median and interquartile range (IQR) and differences between groups were first analysed using Kruskal-Wallis test and then by Mann-Whitney U test if any significance were found. Correlations be-

tween variables were examined by Spearman rank correlation coefficient (R_s). In order to reduce the possibility of correlations occurring by chance, only those with $R_s > 0.50$ and significant at ≤ 0.01 level were considered relevant. All other tests considered a p value of ≤ 0.05 as statistically significant. All data were analysed using the statistical package GraphPad Prism® version 2.01 for Window 95 and NT.

RESULTS

Sixteen CS-dependant asthmatics (7 on high dose inhaled CS alone and 9 on additional oral CS), 12 mild asthmatics and 10 normal subjects were recruited. The clinico-demographic characteristics of these subjects are described in Table 1. Generally, CS-dependant asthmatics were older, had asthma longer and demonstrated significantly lower FEV₁ than mild stable asthmatics. Between the two groups of CS-dependant asthmatics, those requiring additional oral CS showed a trend towards older age, longer disease duration and lower FEV₁.

The maximal percentage FEV₁ fall following nebulized hypertonic saline was significantly greater in CS-dependent asthmatics than in mild asthmatics or normal subjects (on inhaled CS: median [IQR]: 16.0 [12.3-22.0]; on oral CS: 16.0 [11.0-21.0]; mild asthmatics, 5.3 [4.2-10.8]; normal subjects: 4.6 [3.4-6.4]) (Fig. 1). Between those asthmatics dependent on inhaled CS and oral CS, the FEV₁ fall was comparable. Similarly, there was no significant difference in FEV₁ fall between mild asthmatics and normal subjects. The inverse correlation between maximal percentage FEV₁ fall induced by the procedure and baseline clinic FEV₁ in all asthmatics was highly significant ($R_s = -0.69$; $p < 0.0001$) (Fig. 2).

Two inhaled CS-dependant asthmatics, 4 oral CS-dependant asthmatics and 1 mild asthmatics had FEV₁ fall $\geq 20\%$. However, among them, only 3 oral CS-dependant and 1 inhaled CS-dependant asthmatics had a clinically significant event with marked wheezing and chest tightness. Their bronchospasm were easily reversed with nebulized β_2 -agonist within 30 minutes and did not required prolonged observation or hospitalisation. Three of these

Table 1 Clinico-demographic characteristics of asthmatic and normal subjects

Characteristics	CS-dependant asthmatics			Mild stable asthmatics	Healthy subjects
	Whole group	On high dose inhaled CS	On additional oral CS		
N	16	7	9	12	10
Age (years)	48** (30-60)	42 (30-55)	51 (42-60)	25 (23-28)	26 (23-30)
Male, N	11	6	5	9	6
FEV₁ % predicted normal	46.3** (27.4-69.4)	48.4 (27.4-69.4)	43.9 (31.2-56.6)	93.6* (86.0-101.1)	107.1 (98.8-115.4)
FEV₁ liters/minute	1.45** (0.9-2.5)	1.82 (1.2-2.5)	1.33 (0.9-1.8)	3.95 (3.4-4.5)	4.18 (3.5-4.9)
Disease duration (years)	31** (9-44)	21 (9-34)	33 (23-44)	16 (13-20)	-
Daily CS requirement	-	1,150 (800- 1,600) µg [†]	18.4 (9- 40) mg [‡]	none	-
Other controller medications					
Theophylline	13	5	8	none	-
Salmeterol	16	7	9	none	-
Ipratropium bromide	13	4	9	none	-
Immunosuppressants[¶]	3	0	3	none	-

Values are mean (range) unless otherwise specified

CS= corticosteroids; FEV₁= Forced Expiratory Volume in One Second;

[¶]Intended as corticosteroid-sparing agents. All three patients previously had cyclosporin A

[†]inhaled budesonide or equivalent; [‡] prednisolone

p* < 0.05 versus normal subjects; *p* < 0.01 versus mild asthmatics or normal subjects






Fig. 1 Maximal percentage FEV₁ fall from baseline following sputum induction. ICS = inhaled CS-dependent asthmatics; OCS = oral CS-dependent asthmatics; Mild = mild asthmatics; Normal = normal subjects. Horizontal bars represent median. ** *p* < 0.01 OCS versus mild and normal and ICS versus normal. **p* < 0.05 ICS versus mild asthmatics.



asthmatics had FEV₁ of < 1 liter. Of the 38 subjects studied, 1 normal subject, 1 mild asthmatic and 3 oral CS-dependant asthmatics could not yield sufficient sputum for processing.

In mild asthmatics, percentage sputum eosinophil was higher than in inhaled CS-dependent asthmatics (median [IQR] %: 23.3 [10.7-46.3] versus 2.3 [1.6-4.6], $p = 0.006$) and in oral CS-dependent asthmatics (versus 4.7 [1.0-12.6], $p = 0.05$). Percentage sputum neutrophil in CS-dependent asthmatics (on inhaled CS, 68.9 [51.5-78.9]; on oral CS, 62.1 [54.6-71.9]) was significantly higher than those in mild asthmatics (28.7 [19.0-42.6], $p = 0.03$ and 0.04, respectively), but comparable to those in healthy subjects (44.7 [30.3-49.5]). Percentage sputum macrophages and lymphocytes were comparable between these groups (Fig. 3).

Percentage sputum eosinophil did not significantly correlate to baseline FEV₁ in either CS-dependent ($R_s = -0.26$, $p = 0.39$), non-CS-requiring ($R_s = -0.17$, $p = 0.61$) or when all asthmatics were studied together ($R_s = 0.38$, $p = 0.06$) (Fig. 4, top). Similarly, there was no significant correlation between percentage sputum neutrophils and baseline FEV₁ in either CS-dependent ($R_s = -0.19$, $p = 0.53$) or non-CS-requiring ($R_s = -0.15$, $p = 0.65$) asthmatics alone. When all asthmatics were studied together, the inverse correlation between percentage sputum neutrophils and FEV₁ was statistically significant ($R_s = -0.51$, $p = 0.01$) (Fig. 4, bottom). In other sputum cell types, there was no significant correlation between percentage counts with baseline FEV₁ (data not shown).

DISCUSSIONS

We have shown that sputum induction using nebulized hypertonic saline can induce significant bronchoconstriction in symptomatic CS-dependant asthmatics, when compared to mild stable asthmatics and normal healthy subjects. The maximal percentage FEV₁ fall correlated inversely with baseline lung function of these asthmatics. Significant bronchoconstriction still occurred despite pre-treatment with short-acting β_2 -agonist. However, they were easily reversed with nebulized short-acting β_2 -agonist and there were no subjects that required prolonged observation or hospitalization. Induction of sufficient




Fig. 4 Correlations between FEV₁ and percentage sputum eosinophils (TOP) and neutrophils (bottom) in asthmatics. Mild asthmatics (◆); inhaled CS-dependant asthmatics (○); oral CS-dependant asthmatics (▲).

sputum for analysis using this method was generally successful in CS-dependant patients.

We also showed that the proportion of sputum eosinophils and neutrophils in induced sputum differed appreciably between CS-dependant and mild stable asthmatics. Eosinophils were significantly lower, while neutrophils, significantly higher in CS-dependant subjects. The proportion of sputum neutrophils in CS-dependant asthmatics were however comparable to those from healthy subjects and furthermore, there was considerable overlap between all study populations, suggesting that these were highly variable between subjects. While there was no significant correlation between sputum eosinophils and FEV₁, a significant inverse correlation could be demonstrated between percentage sputum neutrophils and FEV₁ when all asthmatics were studied together.

Over the last few years, the modern method of sputum induction and processing as an approach to study airway inflammation has gained wide acceptance. Many studies have now shown that the induced sputum obtained from this approach are repeatable, reliable¹⁻³ and responsive to change.^{4,16} The obvious advantage in the use of induced sputum is that it ameliorates the risk associated with bronchoscopy when studying subjects with poor or unstable lung function.

Two studies have specifically addressed the safety issue of performing sputum induction using nebulized hypertonic saline in asthmatic patients. In 64 asthmatics of varying severity and dependency of CS therapy, de la Fuente *et al.*⁵ showed that severe asthmatics had a significantly greater FEV₁ fall following this procedure than mild asthmatics. However, the FEV₁ fall in their subjects did not exceed 20%. Furthermore, they could not show any significant correlation between FEV₁ fall and baseline FEV₁. In another study by Grootendorst *et al.*⁶ of 20 adolescent asthmatics with moderate-to-severe severity dependent on high dose inhaled CS, sputum induction with hypertonic saline brought about an increase of about 9% FEV₁ predicted, suggesting that there was very little risk associated with the procedure in their asthmatic patients.

Our findings add to the bulk of existing data on safety of inducing sputum with nebulized hypertonic saline in that for symptomatic asthmatics despite being treated with high doses of inhaled CS with or without additional oral CS, sputum induction with nebulized hypertonic saline represents a definite risk. In these patients, the resulted FEV₁ fall can exceed 20% and clinically significant wheezing can occur. Furthermore, baseline FEV₁ can predict a risk of significant bronchoconstriction. One likely explanation for the study of de la Fuente *et al.*⁵ not being able to show significant FEV₁ falls is that they had intentionally excluded asthmatics with baseline FEV₁ < 1 liter in their study population. This is relevant because three of our four asthmatics who developed clinically significant wheeze had baseline FEV₁ < 1 liter.

Currently there is a move towards using isotonic saline, instead of hypertonic saline to induce sputum in patients with poor or unstable lung func-

tion. This is intuitively appropriate since it has been well recognized that inhalation of hypertonic saline can cause bronchoconstriction in normal and asthmatic subjects.^{17,18} Sputum cellular profile in asthmatics has been shown to be slightly altered whether they were induced using hypertonic or isotonic saline.²⁰ Furthermore, isotonic saline has been shown to safely induced sputum in children²⁰ and adults⁴ during asthma exacerbation.

Our findings on sputum cellular profile support the view that in some asthmatics chronically dependent on CS, neutrophils may dominate in the airways.^{11,7} Wenzel *et al.*¹¹ showed that in the bronchial mucosa of a group of symptomatic CS-dependant asthmatics not dissimilar to ours, the number of neutrophils in both bronchial biopsies and bronchoalveolar lavage (BAL) was significantly higher than in moderate asthmatics, not on CS treatment and in normal subjects. Similarly, Jatakanon *et al.*⁷ showed that in induced sputum, neutrophil numbers were significantly higher in severe CS-dependant asthmatics than in mild asthmatics or in normal subjects. This view that CS-dependant asthmatics have more airway neutrophilia is however still being debated as there are also other studies showing that sputum eosinophils were increased in severe CS-dependant asthmatics.^{21,22} More recently, studies have suggested asthmatic patients with neutrophilic airway inflammation may represent a subgroup of asthmatics that is less responsive to the effect of corticosteroids.^{23,24} Another possible perspective that is worth considering is whether chronic CS use in asthma is responsible for this alternation of airway 'phenotype'. It has been recognized that while CS could increase eosinophil apoptosis,^{25,26} it could paradoxically delay neutrophil apoptosis²⁶ and therefore lead to a dominance of airway neutrophils. Whether such neutrophils play any pathogenic role under such circumstances is unclear and warrants further research.

ACKNOWLEDGMENTS

The authors wish to thank the support of staff in chest and research clinics in the identification and recruitment of patients and normal volunteers. The study was funded by GlaxoSmithKline Ltd and internal university research fund.

REFERENCES

1. Pizzichini E, Pizzichini M, Efthimiadis A, *et al.* Indices of airway inflammation in induced sputum: reproducibility and validity of cell and fluid-phase measurements. *Am J Respir Crit Care Med* 1996; 154: 308-17.
2. Spanevello A, Migliori GB, Sharara A, *et al.* Induced sputum to assess airway inflammation: a study of reproducibility. *Clin Exp Allergy* 1997; 27: 1138-44.
3. in 't Veen JC, de Gouw HW, Smits HH, *et al.* Repeatability of cellular and soluble markers of inflammation in induced sputum from patients with asthma. *Eur Respir J* 1996; 9: 2441-7.
4. Pizzichini MM, Pizzichini E, Clelland L, *et al.* Sputum in severe exacerbations of asthma: kinetics of inflammatory indices after prednisone treatment. *Am J Respir Crit Care Med* 1997; 155: 1501-8.
5. de la Fuente PT, Romagnoli M, Godard P, Bousquet J, Chanez P. Safety of inducing sputum in patients with asthma of varying severity. *Am J Respir Crit Care Med* 1998; 157: 1127-30.
6. Grootendorst DC, van den Bos JW, Romeijn JJ, *et al.* Induced sputum in adolescents with severe stable asthma. Safety and the relationship of cell counts and eosinophil cationic protein to clinical severity. *Eur Respir J* 1999; 13: 647-53.
7. Jatakanon A, Uasuf C, Maziak W, Lim S, Chung KF, Barnes PJ. Neutrophilic inflammation in severe persistent asthma. *Am J Respir Crit Care Med* 1999; 160: 1532-9.
8. Bousquet J, Chanez P, Lacoste JY, *et al.* Eosinophilic inflammation in asthma. *N Engl J Med* 1990; 323: 1033-9.
9. Ronchi MC, Piragino C, Rosi E, *et al.* Do sputum eosinophils and ECP relate to the severity of asthma? *Eur Respir J* 1997; 10: 1809-13.
10. Fujimoto K, Kubo K, Matsuzawa Y, Sekiguchi M. Eosinophil cationic protein levels in induced sputum correlate with the severity of bronchial asthma. *Chest* 1997; 112: 1241-7.
11. Wenzel SE, Szefler SJ, Leung DY, Sloan SI, Rex MR, Martin RJ. Bronchoscopic evaluation of severe asthma. *Am J Respir Crit Care Med* 1997; 156: 737-43.
12. International consensus report on diagnosis and treatment of asthma. National Heart, Lung, and Blood Institute, National Institutes of Health. Bethesda, Maryland 20892. Publication no. 92-3091, March 1992.
13. Tattersfield AE, Holmes W. Who should look after asthma? *Thorax* 1995; 50: 597-9.
14. British Thoracic Society Guidelines on the management of asthma. *Thorax* 1993; 48: S1-S24.
15. Pin I, Gibson PG, Kolendowicz R, *et al.* Use of induced sputum cell counts to investigate airway inflammation. *Thorax* 1992; 47: 25-9.
16. Fahy JV, Liu J, Wong H, Boushey HA. Analysis of cellular and biochemical constituents of induced sputum after allergen challenge: a method for studying allergic airway inflammation. *J Allergy Clin Immunol* 1994; 93: 1031-9.
17. Sont JK, Booms P, Bel EH, Vandembroucke JP, Sterk PJ. The severity of breathlessness during challenges with inhaled methacholine and hypertonic saline in atopic asthmatic subjects. The relationship with deep breath-induced bronchodilation. *Am J Respir Crit Care Med* 1995; 152: 38-44.
18. Makker HK, Holgate ST. Relation of the hypertonic saline responsiveness of the airways to exercise induced asthma symptom severity and to histamine or methacholine reactivity. *Thorax* 1993; 48: 142-7.
19. Bacci E, Cianchetti S, Paggiaro PL, *et al.* Comparison between hypertonic and isotonic saline-induced sputum in the evaluation of airway inflammation in subjects with moderate asthma. *Clin Exp Allergy* 1996; 26: 1395-1400.
20. Twaddell SH, Gibson PG, Carty K, Woolley KL, Henry RL. Assessment of airway inflammation in children with acute asthma using induced sputum. *Eur Respir J* 1996; 9: 2104-8.
21. Tarodo de la Fuente P, Romagnoli M, Carlsson L, Godard P, Bousquet J, Chanez P. Eosinophilic inflammation assessed by induced sputum in corticosteroid-dependent asthma. *Respir Med* 1999; 93: 183-9.
22. Louis R, Lau LC, Bron AO, Roldaan AC, Radermecker M, Djukanovic R. The relationship between airways inflammation and asthma severity. *Am J Respir Crit Care Med* 2000; 161: 9-16.
23. Green RH, Brightling CE, Woltmann G, Parker D, Wardlaw AJ, Pavord ID. Analysis of induced sputum in adults with asthma: identification of subgroup with isolated sputum neutrophilia and poor response to inhaled corticosteroids. *Thorax* 2002; 57: 875-9.
24. Gibson PG, Simpson JL, Saltos N. Heterogeneity of airway inflammation in persistent asthma: evidence of neutrophilic inflammation and increased sputum interleukin-8. *Chest* 2001; 119: 1299-300.
25. Woolley KL, Gibson PG, Carty K, Wilson AJ, Twaddell SH, Woolley MJ. Eosinophil apoptosis and the resolution of airway inflammation in asthma. *Am J Respir Crit Care Med* 1996; 154: 237-43.
26. Meagher LC, Cousin JM, Seckl JR, Haslett C. Opposing effects of glucocorticoids on the rate of apoptosis in neutrophilic and eosinophilic granulocytes. *J Immunol* 1996; 156: 4422-8.