

# Inhibition of Tryptase Release from Human Colon Mast Cells by Histamine Receptor Antagonists

Shao-Heng He<sup>1,2</sup>, Hua Xie<sup>1</sup> and Yi-Ling Fu<sup>1</sup>

**SUMMARY** The main objective of this study was to investigate the ability of histamine receptor antagonists to modulate tryptase release from human colon mast cells induced by histamine. Enzymatically dispersed cells from human colon were challenged with histamine in the absence or presence of the histamine receptor antagonists, and the tryptase release was determined. It was found that histamine induced tryptase release from colon mast cells was inhibited by up to approximately 61.5% and 24% by the H<sub>1</sub> histamine receptor antagonist terfenadine and the H<sub>2</sub> histamine receptor antagonist cimetidine, respectively, when histamine and its antagonists were added to cells at the same time. The H<sub>3</sub> histamine receptor antagonist clobenpropit had no effect on histamine induced tryptase release from colon mast cells at all concentrations tested. Preincubation of terfenadine, cimetidine or clobenpropit with cells for 20 minutes before challenging with histamine did not enhance the ability of these anti-histamines to inhibit histamine induced tryptase release. Apart from terfenadine at 100 µg/ml, the antagonists themselves did not stimulate tryptase release from colon mast cells following both 15 minutes and 35 minutes incubation periods. It was concluded that H<sub>1</sub> and H<sub>2</sub> histamine receptor antagonists were able to inhibit histamine induced tryptase release from colon mast cells. This not only added some new data to our hypothesis of self-amplification mechanisms of mast cell degranulation, but also suggested that combining these two types of anti-histamine drugs could be useful for the treatment of inflammatory bowel disease (IBD).

It was reported that an increase in mast cells and mast cell degranulation are closely associated to several gastrointestinal diseases including idiopathic inflammatory bowel disease,<sup>1</sup> chronic ulcerative colitis,<sup>2</sup> Crohn's disease,<sup>3-5</sup> gastritis,<sup>5</sup> collagenous colitis,<sup>6,7</sup> irritable bowel syndrome<sup>8,9</sup> and chronic inflammatory duodenal bowel disorders.<sup>10</sup> Through releasing their proinflammatory mediators including histamine, tryptase, chymase, heparin and some cytokines,<sup>11</sup> mast cells actively participate in the pathogenesis of these intestinal diseases.

Histamine has been found to have a close relationship with IBD. For example, the secretion rate of histamine was increased in patients with active

Crohn's disease compared to normal controls, and the secretion of histamine was related to the disease activity;<sup>12</sup> highly elevated mucosal histamine levels were observed with allergic enteropathy and ulcerative colitis<sup>13</sup> and an enhanced histamine metabolism was found with collagenous colitis and food allergy.<sup>14</sup>

Histamine is a primary amine synthesized from histidine in the Golgi apparatus, from where it is transported to the granules for storage in ionic associa-

From the <sup>1</sup>Allergy & Inflammation Research Institute, Shantou University Medical College, Shantou 515031, China, <sup>2</sup>Immunopharmacology Group, University of Southampton, Southampton, U.K.

Correspondence: Shao-Heng He  
E-mail: shoahenghe@hotmail.com

tion with the acidic residues of the glycosaminoglycan side chains of heparin and with proteinases. As only mast cells contain histamine in man (apart from histaminergic nerves and a few basophils in human tissues), histamine can be used as a marker for mast cell degranulation.

It was found recently that histamine was able to stimulate tryptase release from human colon mast cells,<sup>15</sup> which may represent a unique self-amplification mechanism of mast cell degranulation. However, the mechanism by which histamine acts on mast cells remains unclear. To date, a total of three histamine receptors H<sub>1</sub>, H<sub>2</sub>, and H<sub>3</sub> have been discovered in the human gut,<sup>16,17</sup> and they are all expressed on human mast cells.<sup>18</sup> This suggests that histamine may act on mast cells through these receptors. Since tryptase is a unique marker of mast cell degranulation,<sup>19</sup> we investigated the effects of histamine receptor antagonists on histamine induced tryptase release from human colon mast cells in the current study.

## MATERIALS AND METHODS

### Dispersion of mast cells

Human colon tissue was obtained from patients with carcinoma of the colon at colectomy. Only macroscopically normal tissue was used for the study. After removing fat, the tissue was washed and chopped finely with scissors into fragments of 0.5-2.0 mm<sup>3</sup>, and then incubated with 1.5 mg/ml collagenase (Sigma) and 0.75 mg/ml hyaluronidase (Sigma) in MEM containing 2% fetal calf serum (FCS; 1 g colon/10 ml buffer) for 70 minutes at 37°C. Dispersed cells were separated from undigested tissue by filtration through nylon gauze (pore size 100 µm diameter), washed and then maintained in MEM (Gibco) (containing 10% FCS, 200 U/ml penicillin, 200 µg/ml streptomycin) on a roller overnight at room temperature. Mast cell purity, as determined by light microscopy after staining with Alcine blue, ranged from 3.5 to 5.4%.

### Mast cell challenge

Dispersed cells were resuspended in HEPES buffered salt solution (HBSS, pH 7.4) with CaCl<sub>2</sub> and MgCl<sub>2</sub> (complete HBSS), and 100 µl aliquots containing 4-6 x 10<sup>3</sup> mast cells were added to 50 µl histamine (Sigma), the calcium ionophore A23187

(CI, Sigma), or a histamine receptor antagonist in complete HBSS and incubated for 15 minutes or 35 minutes at 37°C, respectively. The reaction was terminated by addition of 150 µl ice cold incomplete HBSS and the tubes were centrifuged immediately (500 x g, 10 minutes, 4°C). All experiments were performed in duplicate. Supernatants were stored at -20°C until tryptase concentrations were determined.

### Inhibition of the tryptase release

For some experiments, the histamine receptor antagonist was preincubated with the cells for 20 minutes before histamine was added. For other experiments, the histamine receptor antagonist and histamine were added to the cells at the same time (no preincubation period). Histamine and the antagonist were then incubated with the cells at 37°C for 15 minutes. Data were expressed as the percentage inhibition of tryptase release, taking into account tryptase release in the presence and absence of the anti-histamine. Based on our previous experiments, histamine concentrations at 100 µg/ml and 1,000 µg/ml were taken as the standard concentrations throughout the study.<sup>15</sup>

### Tryptase measurement

Tryptase concentrations were measured with a sandwich ELISA procedure with a specific polyclonal antibody against human tryptase as the capture antibody and AA5, a monoclonal antibody specific for human tryptase as the detecting antibody.<sup>20</sup>

### Statistical analyses

Statistical analyses were performed with SPSS software. Data were expressed as mean ± SEM. Where analysis of variance indicated significant differences between groups with ANOVA, for the pre-planned comparisons of interest, Student's t test was applied. For all analyses, *p* < 0.05 was taken as significant.

## RESULTS

### Effects of secretagogues and anti-histamine on tryptase release from mast cells

At 15 minutes following incubation, histamine at 100 µg/ml and 1,000 µg/ml and CI at 1 µg/ml were able to induce 25.7 ± 5.8, 42.4 ± 13.8 and 27.9 ±

3.6 ng/ml tryptase release from colon mast cells respectively, whereas at the same time point spontaneous tryptase release (buffer alone) was  $19.4 \pm 4.7$  ng/ml. Following a 35 minutes incubation period, 100  $\mu\text{g/ml}$  and 1,000  $\mu\text{g/ml}$  histamine were able to provoke  $23.8 \pm 5.1$  and  $26.7 \pm 5.1$  ng/ml tryptase release, respectively. The  $H_1$  histamine receptor antagonist terfenadine at 100  $\mu\text{g/ml}$  induced significant tryptase release following both 15 minutes and 35 minutes incubation periods. The  $H_2$  histamine receptor antagonist cimetidine and the  $H_3$  histamine receptor antagonist clobenpropit at all concentrations tested had no stimulatory effect on colon mast cells (Table 1).

#### Effects of anti-histamines on histamine induced tryptase release from mast cells

The concentration dependent inhibition of histamine induced release of tryptase from colon mast cells was observed when 1,000  $\mu\text{g/ml}$  histamine and various concentrations of terfenadine were added to the cells at the same time (Fig. 1). Approximately 61.5% inhibition of histamine induced tryptase release was achieved with 100  $\mu\text{g/ml}$  terfenadine (Fig. 1). As little as 1.0  $\mu\text{g/ml}$  terfenadine was able to sig-

nificantly inhibit histamine induced tryptase release when histamine and terfenadine were added to the cells at the same time. Similarly, but to a much lesser extent, cimetidine at a concentration of 100  $\mu\text{g/ml}$  was able to reduce histamine induced tryptase release by approximately 24%. Clobenpropit had little effect on histamine induced tryptase release. Terfenadine and cimetidine were also able to inhibit 100  $\mu\text{g/ml}$  histamine induced tryptase release (data not shown). Preincubation of terfenadine, cimetidine or clobenpropit with cells for 20 minutes before challenging with histamine did not enhance the ability of these anti-histamines to inhibit histamine induced tryptase release (Fig. 2).


#### DISCUSSION

It was interesting to learn that the histamine  $H_1$  receptor antagonist terfenadine was able to abolish more than 60% histamine induced tryptase release from colon mast cells, which indicated strongly that histamine induced tryptase release from colon mast cells were mainly through the  $H_1$  receptor. Approximately 24% histamine induced tryptase release was inhibited by the histamine  $H_2$  receptor antagonist cimetidine implicating that activation of mast cells

**Table 1** Effect of histamine and its receptor antagonists on tryptase release from human colon mast cells


Compound ( $\mu\text{g/ml}$ )	Tryptase released (ng/ml)	
	15 minutes	35 minutes
Buffer alone	$19.4 \pm 4.7$	$17.8 \pm 3.0$
Terfenadine	0.1	$19.5 \pm 4.1$
	1.0	$24.3 \pm 4.9$
	10	$24.2 \pm 5.1$
	100	$26.7 \pm 2.1^a$
Cimetidine	0.1	$24.1 \pm 4.3$
	1.0	$23.3 \pm 3.5$
	10	$22.3 \pm 3.7$
	100	$17.4 \pm 4.0$
Clobenpropit	0.1	$20.3 \pm 3.8$
	1.0	$20.6 \pm 5.1$
	10	$20.0 \pm 4.2$
	100	$23.0 \pm 4.9$
Histamine	100	$25.7 \pm 5.8^a$
	1000	$42.4 \pm 13.8^a$
CI	1.0	$27.9 \pm 3.6^a$

The values are mean  $\pm$  SEM for five separate experiments.  $a$   $p < 0.05$  compared with buffer alone control (Student's  $t$  test). CI = calcium ionophore.



by histamine was partially through  $H_2$  receptors. The phenomenon that histamine acts on both  $H_1$  and  $H_2$  receptors for one pathophysiological event was observed previously when a pyrilamine and cimetidine combination was able to completely abolish histamine induced microvascular leakage in a guinea pig model.<sup>21</sup> In contrast to  $H_1$  and  $H_2$  receptors,  $H_3$  receptors seemed to play no role in histamine induced activation of mast cells, as its antagonist clobenpropit did not show any effect on histamine induced tryptase release.

Cimetidine by itself did not stimulate tryptase release from mast cells indicating that within the doses tested cimetidine may be safe to be used as a mast cell stabilizer, whereas the dose of terfenadine should not exceed 10 µg/ml. Since the histamine content of mast cells is 2 to 5 pg/cell, 1,000 µg/ml histamine could only be achieved at the nearby surrounding area of the degranulating mast cells, which indirectly suggested that the potent inhibition of his-



tamine induced tryptase release by terfenadine should efficiently block histamine induced mast cell degranulation. The rapid blockage of histamine induced mast cell activation by anti-histamine drugs and the close relationship between histamine and IBD suggested that a combination of  $H_1$  and  $H_2$  histamine receptor antagonists might be useful for the treatment of IBD.

The evidence that actions of histamine on mast cells was through  $H_1$  and  $H_2$  histamine receptors added some new data in our hypothesis of self-amplification mechanisms of mast cell degranulation,<sup>22</sup> which is that degranulated mast cells release histamine, histamine then either acts on its surrounding mast cells (paracrine mechanism) or back on its host mast cells (autocrine mechanism) through  $H_1$  and  $H_2$  histamine receptors.

In conclusion, the H<sub>1</sub> and H<sub>2</sub> histamine receptor antagonists were able to inhibit histamine induced tryptase release from colon mast cells, indicating that combining these two types of anti-histamine drugs may be useful for the treatment of IBD.

### ACKNOWLEDGEMENTS

This project was supported by the Major State Basic Research Program of China (973 Program), China (No. 2001CB510009); the Planned Science and Technology Project of Guangdong Province, China (No. 2003B31502) and the Li Ka Shing Foundation, Hong Kong, China (No. C0200001).

### REFERENCES

1. Fox CC, Lichtenstein LM, Roche JK. Intestinal mast cell responses in idiopathic inflammatory bowel disease. Histamine release from human intestinal mast cells in response to gut epithelial proteins. *Dig Dis Sci* 1993; 38: 1105-12.
2. Stoyanova II, Gulubova MV. Mast cells and inflammatory mediators in chronic ulcerative colitis. *Acta Histochem* 2002; 104: 185-92.
3. Gelbmann CM, Mestermann S, Gross V, Kollinger M, Scholmerich J, Falk W. Structures in Crohn's disease are characterised by an accumulation of mast cells colocalised with laminin but not with fibronectin or vitronectin. *Gut* 1999; 45: 210-7.
4. Bischoff SC, Wedemeyer J, Herrmann A, Meier PN, Trautwein C, Cetin Y, Maschek H, Stolte M, Gebel M, Manns MP. Quantitative assessment of intestinal eosinophils and mast cells in inflammatory bowel disease. *Histopathology* 1996; 28: 1-13.
5. Beil WJ, Schulz M, McEuen AR, Buckley MG, Walls AF. Number, fixation properties, dye-binding and protease expression of duodenal mast cells: comparisons between healthy subjects and patients with gastritis or Crohn's disease. *Histochem J* 1997; 29: 759-73.
6. Nishida Y, Murase K, Isomoto H, Furusu H, Mizuta Y, Riddell RH, Kohno S. Different distribution of mast cells and macrophages in colonic mucosa of patients with collagenous colitis and inflammatory bowel disease. *Hepato-gastroenterology* 2002; 49: 678-82.
7. Schwab D, Raithel M, Hahn EG. Evidence for mast cell activation in collagenous colitis. *Inflamm Res* 1998; 47 Suppl 1: S64-5.
8. O'Sullivan M, Clayton N, Breslin NP, Harman I, Bountra C, McLaren A, O'Morain CA. Increased mast cells in the irritable bowel syndrome. *Neurogastroenterol Motil* 2000; 12: 449-57.
9. Pang X, Boucher W, Triadafilopoulos G, Sant GR, Theoharides TC. Mast cell and substance P-positive nerve involvement in a patient with both irritable bowel syndrome and interstitial cystitis. *Urology* 1996; 47: 436-8.
10. Crivellato E, Finato N, Isola M, Ribatti D, Beltrami CA. Low mast cell density in the human duodenal mucosa from chronic inflammatory duodenal bowel disorders is associated with defective villous architecture. *Eur J Clin Invest* 2003; 33: 601-10.
11. Walls AF, He S, Buckley MG, McEuen AR. Roles of the mast cell and basophil in asthma. *Clin Exp Allergy Rev* 2001; 1: 68-72.
12. Knutson L, Ahrenstedt O, Odland B, Hallgren R. The jejunal secretion of histamine is increased in active Crohn's disease. *Gastroenterology* 1990; 98: 849-54.
13. Raithel M, Matek M, Baenkler HW, Jorde W, Hahn EG. Mucosal histamine content and histamine secretion in Crohn's disease, ulcerative colitis and allergic enteropathy. *Int Arch Allergy Immunol* 1995; 108: 127-33.
14. Schwab D, Hahn EG, Raithel M. Enhanced histamine metabolism: a comparative analysis of collagenous colitis and food allergy with respect to the role of diet and NSAID use. *Inflamm Res* 2003; 52: 142-7.
15. He S, Xie H. Modulation of tryptase secretion from human colon mast cells by histamine. *World J Gastroenterol* 2004; 10: 323-6.
16. Bertaccini G, Coruzzi G. An update on histamine H<sub>3</sub> receptors and gastrointestinal functions. *Dig Dis Sci* 1995; 40: 2052-63.
17. Rangachari PK. Histamine: mercurial messenger in the gut. *Am J Physiol* 1992; 262: G1-13.
18. Bent S, Fehling U, Braam U, Schunack W, Schmutzler W. The influence of H<sub>1</sub>-, H<sub>2</sub>- and H<sub>3</sub>-receptors on the spontaneous and con A induced histamine release from human adenoidal mast cells. *Agents Actions* 1991; 33: 67-70.
19. Walls AF. Structure and function of human mast cell tryptase. In: Marone G, ed. *Mast cells and basophils*. New York: Academic Press; 2000; pp. 291-309.
20. Buckley MG, Walters C, Wong WM, Cawley MID, Ren S, Schwartz LB, Walls AF. Mast cell activation in arthritis: detection of  $\alpha$ - and  $\beta$ -tryptase, histamine and eosinophil cationic protein in synovial fluid. *Clin Sci* 1997; 93: 363-70.
21. He S, Walls AF. Human mast cell tryptase: a stimulus of microvascular leakage and mast cell activation. *Eur J Pharmacol* 1997; 328: 89-97.
22. He S. The key role of mast cells and their major secretory products in inflammatory bowel disease. *World J Gastroenterol* 2004; 10: 309-18.